

AMPHIBIA: ANURA: LEPTODACTYLIDAE

Leptodactylus natalensis

Catalogue of American Amphibians and Reptiles.

Heyer, W.R. and M.M. Heyer. 2006. *Leptodactylus natalensis*.

***Leptodactylus natalensis* A. Lutz**
Bubbling Frog

Leptodactylus caliginosus: A. Lutz 1930:2.

Leptodactylus natalensis A. Lutz 1930:7. Type locality, Natal, Brazil (see **Remarks**). Lectotype, National Museum of Natural History (USNM) 81130, adult male, collector and date unknown (see **Remarks**) (examined by WRH).

Leptodactylus petersi (sic, *petersii*): Parker 1933:7 (in part).

Leptodactylus podicipinus: Cochran 1955:326 (in part).

Leptodactylus caliginosus caliginosus: B. Lutz 1954: 177.

Leptodactylus wagneri: Heyer 1970:18 (in part).

• **CONTENT.** The species is monotypic.

• **DEFINITION.** Adult *Leptodactylus natalensis* are of moderately small size, the head is about as wide as long, and the limbs are moderate in length (see Table 1; Heyer and Thompson 2000 provided definitions of adult size and leg length categories for *Leptodactylus*). Male vocal sacs usually are not visible externally or weak and median. Male forearms are not noticeably hypertrophied. Adult males have two keratinized spines on each thumb; males lack chest spines. A single pair of dorsolateral folds is rarely absent (2%), usually short (46%), or moderate (53%) in length. The toe tips are narrow (13%), just swollen (31%), swollen (28%), just expanded (22%), or expanded (6%). The toes are fringed; the fringing unites to form basal webbing between the toes. The upper shank has scattered moderate-sized keratin-tipped tubercles (tubercles are white if keratin tips have fallen off). The outer tarsus has scattered small keratin-tipped tubercles. The sole of the foot is either smooth or has a few small keratin-tipped tubercles.

The upper lip pattern ranges from uniform to three distinct light vertical stripes, one obliquely sloping forward stripe extends from lower loreal region to the jaw, ending just before the level of the nostril. A second, weakly obliquely sloping backward stripe extends from under the anterior portion of eye to the jaw. A third stripe extends from the lower posterior portion of the eye to below the tympanum, and may be either interrupted or continuous with a light commissural gland stripe; light stripes may be dark bordered or not. A dark interorbital triangle is usually bordered at least anteriorly by a white stripe. There are usually two ill-defined middorsal dark chevrons, one in the area above the arm insertion and the second over the sacrum. The dorsolateral folds are somewhat darker than the adjacent areas. The species lacks light middorsal stripes. The belly rarely lacks a pattern (1%), occasionally is lightly mottled (7%), and



Figure 1. *Leptodactylus natalensis*, C.M. Carvalho field number 783 to be deposited in the MZUSP, from the campus of the Universidade Federal de Sergipe, São Cristóvão, Brazil (photograph courtesy of Celso Morato de Carvalho).

usually is either moderately (52%) or extensively (40%) mottled. The posterior thigh surfaces rarely (5%) have distinct light stripes, sometimes (16%) the stripes are indistinct, but usually (79%) light stripes are not discernible. The upper shanks have indistinct to distinct somewhat irregular, dark, moderately broad crossbands.





Figure 2. Tadpole of *Leptodactylus natalensis*, redrawn from figure 3 in Oliveira and Lírio Júnior (2000). Specimen from campus of Universidade Federal de Sergipe, São Cristóvão, Brazil. Upper scale bar = 5 mm, lower scale bar = 0.2 mm.

Larvae have a typical pond morphology and are members of the exotrophic, lentic, benthic guild (McDiarmid and Altig 1999). The oral disk is anteroventrally positioned, entire (not emarginated), with an anterior gap lacking marginal papillae. There is a double row of marginal papillae. No submarginal papillae are present. The tooth row formula is 2/3. The spiracle is sinistral and the vent tube is median. The dorsal fin originates at the body/tail junction. At Gosner stages 39/40 the total length is 28 mm with a body length of about 14 mm; the eye diameter is

about 10% the body length; and the oral disk is about 50% of the body length. The body and tail musculature are ash-brown with gold punctuations; the dorsal and ventral tail fins are uniformly black (description based on Oliveira and Lírio Júnior 2000).

Table 1. Summary measurement data for *Leptodactylus natalensis* (means are in parentheses).

Measurement	Males	Females
SVL (mm)	29–42 (34.5)	33–49 (40.0)
Head length/ SVL (%)	35–42 (38)	33–40 (37)
Head width/ SVL (%)	33–39 (36)	30–38 (34)
Thigh length/ SVL (%)	39–49 (44)	38–48 (43)
Shank length/ SVL (%)	41–48 (46)	41–51 (45)
Foot length/ SVL (%)	44–58 (53)	46–57 (52)

The advertisement call consists of a single note per call, given at a rate of 180–250/min. Call duration ranges from 0.06–0.07s. The call is pulsed or partially pulsed (= incompletely amplitude modulated), with 2–7 pulses or partial pulses per call. The call attains maximum intensity in the first 1/5 to 1/2 of the call and attenuates more or less regularly for the remainder of the call. The calls are frequency modulated with extremely rapid rise times. Beginning carrier (also equal fundamental) frequencies range from averages of 550–770 or 1020–1040 Hz to maximum average carrier frequencies of 1370–1830 Hz. Harmonics are either weak or not discernible.




Figure 3. Wave form and audiospectrogram displays of the advertisement call of *Leptodactylus natalensis* (USNM recording 323, cut 1) from the campus of the Universidade Federal de Sergipe, São Cristóvão, Sergipe, Brazil, voucher specimen MZUSP 89945. The wave form is that of the first call shown on the audiospectrogram.

- **DIAGNOSIS.** The species having distinct toe fringes and no more than a single pair of moderate length dorsolateral folds in some or all individuals are *Leptodactylus colombiensis*, *discodactylus*, *diedrus*, *griseigularis*, *leptodactyloides*, *melanonotus*, *natalensis*, *nesiotus*, *pascoensis*, *petersii*, *podicipinus*, *pusillus*, *sabanensis*, *silvanimbus*, and *wagneri*. *Leptodactylus diedrus* and *discodactylus* have at least one transverse groove on the dorsal surface of the largest toe disk; the toes of *L. natalensis* are not disked or grooved. *Leptodactylus natalensis* (males ≤ 42 mm SVL, females ≤ 49 mm SVL) are smaller than almost all *L. pascoensis* and *wagneri* (males ≥ 39 mm SVL, females ≥ 52 mm SVL). *Leptodactylus natalensis* cannot be consistently diagnosed from the remaining species based on morphology. The only other species that occurs in the same general geographic region with *L. natalensis* is *L. podicipinus*. No individuals of *L. natalensis* have distinct light belly spots whereas many *L. podicipinus* individuals do. Just over half of *L. natalensis* specimens have toe tips larger than narrow or just swollen whereas all *L. podicipinus* specimens examined have toe tips that are either narrow or just swollen.

- **DESCRIPTIONS.** A. Lutz's (1930) original description of *Leptodactylus natalensis* included color. His description of *L. caliginosus* in the same paper is of *L. natalensis*, as well. Heyer (1994) provided detailed morphological descriptions of adult females and males. Oliveira and Lírio Júnior (2000) described morphology and color of adult and larval specimens. Accounts of advertisement and chirp calls of *L. natalensis* are in Heyer and Carvalho (2000), and the karyotype is described in Bogart (1974).


- **ILLUSTRATIONS.** Color photographs of adults are in Izecksohn and Carvalho-e-Silva (2001b), Freitas and Silva (2004, 2005), and Oliveira and Lírio Júnior (2000). A. Lutz (1930) provided color illustrations and black-and-white photographs of *L. natalensis*. His color illustrations and black-and-white photographs of *L. caliginosus* in the same paper (A. Lutz 1930) are of *L. natalensis*. Black-and-white photographs of adults are in Bogart (1974) and Izecksohn and Carvalho-e-Silva (2001b), while an illustration of an adult, as *L. caliginosus*, is in B. Lutz (1954). Oliveira and Lírio Júnior (2000) offered figure drawings of a larva and a larval mouth. A camera lucida tracing of the terminal phalanx is found in Heyer (2000). Wave forms and audiospectrograms of both the advertisement and chirp calls are illustrated in Heyer and Carvalho (2000) and a power spectrum of the advertisement call is also pictured. Bogart (1974) displayed a photo of the karyotype and an ideogram of the normalized length of the chromosome.

- **DISTRIBUTION.** *Leptodactylus natalensis* is associated with northern and central portions of the Atlantic Forest Morphoclimatic Domain (Ab'Sáber 1977) from its most northern extent in the State of Rio Grande do Norte to and including the State of Rio de

Janeiro (see Map).

Fabio (1985 as *L. caliginosus*), Heyer and Carvalho (2000), Izecksohn and Carvalho-e-Silva (2001b), and Oliveira and Lírio Júnior (2000) have found the species in forest and at forest edges. Heyer (1994) noted its capture at 50 m elevation.

References to distribution and/or localities in Brazil include Bogart (1974), Bokermann (1966), Cochran (1955, as *L. podicipinus* in part), Duellman (1999), Fabio (1985, as *L. caliginosus*), Heyer (1994), Heyer and Carvalho (2000), Izecksohn and Carvalho-e-Silva (2001a,b), A. Lutz (1930, as both *L. caliginosus* and *L. natalensis*), B. Lutz (1954, as *L. caliginosus* and *L. caliginosus caliginosus*), Myers (1946, as *L. caliginosus*) Oliveira and Lírio Júnior (2000), and Santos et al. (2004).



Map. Distribution of *Leptodactylus natalensis*. The type-locality is indicated by a circle. Dots mark other localities. A dot may represent more than one site. Published locality data used to generate the map should be considered as secondary sources of information, as we did not confirm all specimen localities. The locality coordinate data and sources are available on a spread sheet at <http://learning.richmond.edu/Leptodactylus>.

• FOSSIL RECORD. None.

• PERTINENT LITERATURE. The following literature is listed by topic. The symbol (M) indicates a species mention, and (S) is a secondary source: **bibliographic information and lists** (Liner 1992); **call and call parameters** (Heyer and Carvalho 2000; A. Lutz 1930 as both *L. caliginosus* and *L. natalensis*; B. Lutz 1954 as *L. caliginosus*); **conservation** (Rodrigues 2003, 2004; Young et al. 2004); **ecology, natural history, reproduction** (De la Riva 1993 (M); Fabio 1985 as *L. caliginosus* (M); Heyer and Carvalho

2000; A. Lutz 1930 as both *L. caliginosus* and *L. natalensis*; B. Lutz 1954 as *L. caliginosus*; Oliveira and Lírio Júnior 2000, Rodrigues 2003, Santos et al. 2004); **field guide** (Freitas and Silva 2004, 2005); **faunal lists** (Freitas and Silva 2004; B. Lutz 1954 as *L. caliginosus*; Myers 1946 as *L. caliginosus*; Rodrigues 2004); **faunal lists** (Freitas and Silva 2004, B. Lutz 1954 as *L. caliginosus*; Myers 1946 as *L. caliginosus*; Rodrigues 2004); **habitat** (Heyer and Carvalho 2000; Jim 1980 as *L. caliginosus* (M)); **karyotype** (Bogart 1974; Heyer 1974 (S), 1994 (S); Heyer and Diment 1974; King 1990 (S); Kuramoto 1990 (S); Moreschalchi 1979 (S), Silva et al. 2000 (M)); **keys** (B. Lutz 1954 as *L. caliginosus caliginosus*; Oliveira and Lírio Júnior 2000); **morphology** (Heyer 1994; Heyer and Carvalho 2000; Izecksohn and Carvalho-e-Silva 2001b; Liu 1935a,b; A. Lutz 1930 as both *L. caliginosus* and *L. natalensis*; B. Lutz 1954 as *L. caliginosus*; Oliveira and Lírio Júnior 2000); **nomenclature and taxonomy** (Cei 1980 (M); Heyer 1970, 1974 (M); A. Lutz 1930 as *L. caliginosus*); **parasite host** (Fabio 1985 as *L. caliginosus*; Fabio and Izecksohn 2001); **relationships and systematics** (Bogart 1974, Cei and Erspamer 1965 (M); Gallopin 1962 (M); Heyer and Diment 1974; Savage 2002 (M)); **species accounts** (Heyer 1994; Izecksohn and Carvalho-e-Silva 2001b; A. Lutz 1930 as both *L. caliginosus* and *L. natalensis*; Oliveira and Lírio Júnior 2000); **species comparisons** (Heyer 1994, 2000; Heyer et al. 2002; A. Lutz 1930 as *L. caliginosus*; B. Lutz and Kloss 1952 (M); Oliveira and Lírio Júnior 2000); **species or taxonomic lists** (Gorham 1966 as *L. podicipinus petersii*, 1974 as *L. podicipinus*; Lavilla 1994 (M); Rocha et al. 2004); **type lists** (Cochran 1961).

• NOMENCLATURAL HISTORY. Parker (1933) synonymized *L. natalensis* A. Lutz 1930 with *L. petersi* (sic for *petersii*) (Steindachner 1864). Cochran (1961) listed *L. natalensis* as a synonym of *L. podicipinus* (Cope 1862). Heyer (1970) considered *L. natalensis* to be a synonym of *L. wagneri* Peters 1862. Bogart (1974), based on karyotypic data, proposed that *L. natalensis* and *L. wagneri* represented separate species. Heyer (1994) supported Bogart's proposal based on morphological data.

• REMARKS. "Rã caliginosa" was coined by B. Lutz (1954) as a common name for *L. natalensis* as *L. caliginosus caliginosus*. Izecksohn and Carvalho-e-Silva (2001b) called the frog "Rã borbulhante," or "Bubbling Frog," in reference to its call. Freitas and Silva (2004) used "Caçote" and (2005) "Caçote-borbulhante" for the species.

A. Lutz gave the type-locality as "Natal. Rio Grande do Norte. Rio Baldo e outros lugares" in the Portuguese text (1930:7) and as "Rio Bahú and other places near Natal (Rio Grande do Norte)" in the English text (1930:26). Rio Baldum, 06° 09' S 35° 08' W, is likely the type-locality (P.E. Vanzolini, pers. comm.). A. Lutz described the type material as "vários machos e fêmeas adultos" (1930:7) and "many adults of both sexes" (1930:26). Heyer

(1970:22) designated USNM 81130, a male with the catalogue data Natal, Brazil, collected July 1925, received from A. Lutz (no collector given), as the lectotype of *L. natalensis*. The lectotype was part of A. Lutz's type series, so presumably the lectotype was collected from Rio Baldum. See Heyer (1994:81) for further discussion of the type series.

• **ETYMOLOGY.** The specific name alludes to the general locality where the frog was initially collected.

• **ACKNOWLEDGMENTS.** Celso Morato de Carvalho and Carlos Alberto Gonçalves da Cruz critically reviewed the manuscript. P. E. Vanzolini clarified the type locality for us. The Smithsonian Institution Library staff of the Natural History branch, Claire Catron, Ron Lindsey, Leslie Overstreet, Martha Rosen, Courtney Shaw, David Steere, Wanda West, and Daria Wingreen obtained some of the publications cited in this paper. Research for this account was supported by NSF award DEB-03429 to Rafael O. de Sá and WRH.

LITERATURE CITED

- Ab'Sáber, A.N. 1977. Os domínios morfoclimáticos na América do Sul. Primeira aproximação. *Geomorfologia* (52):1–23 + 1 map.
- Bogart, J.P. 1974. A karyosystematic study of frogs in the genus *Leptodactylus* (Anura: Leptodactylidae). *Copeia* 1974:728–737.
- Bokermann, W.C.A. 1966. Lista Anotada das Localidades Tipo de Anfíbios Brasileiros. Serviço de Documentação - RUSP, São Paulo.
- Cei, J.M. 1980. Amphibians of Argentina. *Monitore Zoologico Italiano* (N.S.), Monografia (2):xii + 609 p.
- and V. Erspamer. 1965. Relaciones taxonómicas y evolución bioquímica de las aminas biógenas en el género *Leptodactylus*. Acta III Congreso Latinoamericano de Zoología, Santiago de Chile. Investigaciones Científicas Zoológicas Chilenas, (separately printed and numbered):1–8.
- Cochran, D.M. 1955 ("1954"). Frogs of Southeastern Brazil. *Bulletin of the United States National Museum* (206):xvi + 423 p.
- . 1961. Type specimens of reptiles and amphibians in the U.S. National Museum. *Bulletin of the United States National Museum* (220):xv + 291.
- De la Riva, I.J. 1993. Ecología de una comunidad neotropical de anfibios durante la estación lluviosa. Tesis Doctoral, Universidad Complutense de Madrid, Madrid.
- Duellman, W.E. 1999. Distribution patterns of amphibians in South America, p. 255–328. In W.E. Duellman (ed.), *Patterns of Distribution of Amphibians. A Global Perspective*. The Johns Hopkins University Press, Baltimore, London.
- Fabio, S.P. 1985 ("1982"). Helmintos de populações simpáticas de algumas espécies de anfíbios anuros da Família Leptodactylidae. Arquivos da Universidade Federal Rural do Rio de Janeiro 5: 69–83.
- and E. Izecksohn. 2001. Ocorrência de *Falcaustra mascula* (Rudolphi, 1819) – (Nematoda: Kathlaniidae) em girino de *Hyla ruschii* Weigoldt [sic] & Peixoto, 1987 (Anura: Hylidae). *Contribuições Avulsas Sobre a História Natural do Brasil* (40):1–6.
- Freitas, M.A. and T.F.S. Silva. 2004. Anfíbios na Bahia. Um Guia de Identificação. Patrocínio Editorial, Políteno, Camaçari, Bahia, Brasil.
- and —. 2005. Guia Ilustrado a Herpetofauna da Mata Atlântica Nordestina. Coleção Manuais de Campo, Editora USEB (União Sul-Americana de Estudos da Biodiversidade), Pelotas, Rio Grande do Sul, Brasil.
- Gallopin, G.C. 1962. El grupo *Pachypus* del género *Leptodactylus* considerado a la luz de las características electroforéticas de sus seroproteínas. *Physis* 23:163–167.
- Gorham, S.W. 1966. Liste der rezenten Amphibien und Reptilien: Ascaphidae, Leiopelmatidea [sic], Pipidae, Discoglossidae, Pelobatidae, Leptodactylidae, Rhinophrynidae. *Das Tierreich* (85):xvi + 222 p.
- . 1974. Checklist of World Amphibians up to January 1, 1970. Liste des Amphibiens du Monde d'après l'État du 1er Janvier 1970. The New Brunswick Museum, Saint John, New Brunswick.
- Heyer, W.R. 1970. Studies on the frogs of the genus *Leptodactylus* (Amphibia: Leptodactylidae). VI. Biosystematics of the *melanonotus* group. *Natural History Museum of Los Angeles County, Contributions in Science* (191):1–48.
- . 1974. Relationships of the *marmoratus* species group (Amphibia, Leptodactylidae) within the Subfamily Leptodactylinae. *Natural History Museum of Los Angeles County, Contributions in Science* (253):1–46.
- . 1994. Variation within the *Leptodactylus podicipinus-wagneri* complex of frogs (Amphibia: Leptodactylidae). *Smithsonian Contributions to Zoology* (546):iv + 1–124.
- . 2000. The lectotype of *Cystignathus hylodes* Reinhardt and Lütken, 1862. *Journal of Herpetology* 34:150–153.
- and C.M. Carvalho. 2000. Calls and calling behavior of the frog *Leptodactylus natalensis* (Amphibia: Anura: Leptodactylidae). *Proceedings of the Biological Society of Washington* 113:284–290.
- , R.O. de Sá, and S. Müller. 2002. *Leptodactylus silvanimbus*. Catalogue of American Amphibians and Reptiles (743):1–3.
- and M.J. Diment. 1974. The karyotype of *Vanzolinia discodactylus* and comments on usefulness of karyotypes in determining relationships in the *Leptodactylus*-complex (Amphibia, Leptodactylidae). *Proceedings of the Biological Society of Washington* 87:327–336.
- and A.S. Thompson. 2000. *Leptodactylus rugosus*. Catalogue of American Amphibians and Reptiles (708):1–5.
- Izecksohn, E. and S.P. Carvalho-e-Silva. 2001a. Anfíbios da Floresta Nacional Mário Xavier, Município

- de Seropédica, Estado do Rio de Janeiro, Brasil (Amphibia: Anura). Contribuições Avulsas sobre a História Natural do Brasil, Série Zoologia (39): 1–3.
- and –. 2001b. Anfíbios do Município do Rio de Janeiro. Editoria UFRJ, Rio de Janeiro.
- Jim, J. 1980. Aspectos Ecológicos dos Anfíbios Registrados na Região de Botucatu, São Paulo (Amphibia, Anura). Tese de Doutorado, Instituto de Biociências, Universidade de São Paulo, São Paulo.
- King, M. 1990. Amphibia, p. vi + 1–241. In J. Bernard and C. Gwent (eds.), Animal Cytogenetics, Volume 4: Chordata 2. Gebrüder Borntraeger, Berlin, Stuttgart.
- Kuramoto, M. 1990. A list of chromosome numbers of anuran amphibians. Bulletin of Fukuoka University of Education 39:83–127.
- Lavilla, E.O. 1994 ("1992"). Tipos portadores de nombre y localidades tipo de anfibios de Argentina. Acta Zoológica Lilloana 42:61–100 +1 unnumbered.
- Liner, E.A. 1992. Bibliography and scientific name index to amphibians and reptiles in the publications of the Biological Society of Washington Bulletin 1–8, 1918–1988 and Proceedings 1–100, 1882–1987. Smithsonian Herpetological Information Service (92):1–68.
- Liu, C.C. 1935a. "The linea masculina," a new secondary sex character in Salientia. Journal of Morphology 57:131–145.
- . 1935b. Types of vocal sac in the Salientia. Proceedings of the Boston Society of Natural History 41:19–40 + plates 4–8.
- Lutz, A. 1930. Segunda memória sobre espécies brasileiras do género *Leptodactylus*, incluindo outras aliadas. Second paper on brasilian and some closely related species of the genus *Leptodactylus*. Memórias do Instituto Oswaldo Cruz 23:1–20 (Portuguese), 21–34 (English) + plates 1–5.
- Lutz, B. 1954. Anfíbios anuros do Distrito Federal. The frogs of the Federal District of Brazil. Memórias do Instituto Oswaldo Cruz 52:155–197 (Portuguese), plates 1–19, 219–238 (English).
- and G.R. Kloss. 1952. Anfíbios anuros do alto Solimões e Rio Negro. Apontamentos sobre algumas formas e suas vicariantes. Short notes on some frogs from the upper Amazonas and a few vicariant forms. Memórias do Instituto Oswaldo Cruz 50: 625–678.
- McDiarmid, R.W. and R. Altig. 1999. Research: Materials and techniques, p. 7–23. In R.W. McDiarmid and R. Altig (eds.), Tadpoles: The Biology of Anuran Larvae. University of Chicago Press, Chicago and London.
- Morescalchi, A. 1979 ("1977"). New developments in vertebrate cytotaxonomy I. Cytotaxonomy of the amphibians. Genetica 50:179–193.
- Myers, G.S. 1946. Lista provisória dos anfíbios do Distrito Federal, Brasil. A provisional list of the amphibians of the Distrito Federal of Brazil. Boletim do Museu Nacional, Nova Série, Zoologia (55):1–18 (Portuguese), 19–36 (English).
- Oliveira, F.F. and G.P. Lírio Júnior. 2000. Anfíbios anuros do campus da Universidade Federal de Sergipe. Biologia Geral e Experimental 1:42–74.
- Parker, H.W. 1933. A list of the frogs and toads of Trinidad. Tropical Agriculture (Trinidad) 10:8–12.
- Rocha, C.F.D., H.G. Bergallo, J.P. Pombal Jr., L. Geise, M. Van Sluys, R. Fernandes, and U. Caramaschi. 2004. Fauna de anfíbios, répteis e mamíferos do estado do Rio de Janeiro, sudeste do Brasil. Publicações Avulsas do Museu Nacional, Rio de Janeiro (104):3–23.
- Rodrigues, M.T. 2003. Herpetofauna da Caatinga, p. 181–236. In I.R. Leal, M. Tabarelli, and J.M.C. Silva (eds.), Ecologia e Conservação da Caatinga. Editora Universitária da Universidade Federal de Pernambuco, Recife, Brasil.
- . 2004. Fauna de anfíbios e répteis das caatingas, p. 173–199. In J.M.C. Silva, M. Tabarelli, M.T. Fonseca, and L.V. Lins (eds.), Biodiversidade da Caatinga: Áreas e Ações Prioritárias para a Conservação. Ministério do Meio Ambiente, Brasília.
- Santos, E.M., A.V. Almeida, and S.D. Vasconcelos. 2004. Feeding habits of six anuran (Amphibia: Anura) species in a rainforest fragment in north-eastern Brazil. Iheringia 94:433–438.
- Savage, J.M. 2002. The Amphibians and Reptiles of Costa Rica. A Herpetofauna between Two Continents, between Two Seas. The University of Chicago Press, Chicago, London.
- Silva, A.P.Z., C.F.B. Haddad, and S. Kasahara. 2000. Chromosomal studies on five species of the genus *Leptodactylus* Fitzinger, 1826 (Amphibia, Anura) using differential staining. Cytobios 103: 25–38.
- Young, B.E., S.N. Stuart, J.S. Chanson, N.A. Cox, and T.M. Boucher. 2004. Disappearing Jewels. The Status of New World Amphibians. NatureServe, Arlington, Virginia.
-
- W. Ronald Heyer and Miriam Muedeking Heyer,** Smithsonian Institution, PO Box 37012, NHB W-201, MRC 162, Washington, DC, 20013–7012, USA (heyerr@si.edu).
- Primary editor for this account, Twan Leenders.
-
- Published 15 February 2006 and Copyright © 2006 by the Society for the Study of Amphibians and Reptiles.