

ECOLOGY

Reprinted from Ecology Vol. 54, No. 3, Late Spring 1973

SPECIES DIVERSITIES OF HERPETOFAUNAL SAMPLES FROM SIMILAR MICROHABITATS AT TWO TROPICAL SITES¹

W. RONALD HEYER AND KEITH A. BERVEN

Biology Department, Pacific Lutheran University, Tacoma, Washington 98447

Abstract. Amphibians and reptiles were collected from tree buttresses in two different regions to determine whether within-habitat species diversity differences were evident. Using the method suggested by Pielou (1966), the average species diversity per individual was $H' = 1.92 \pm .47$ for the collection of amphibians and reptiles taken from tree buttresses in a tropical dry forest formation in Thailand, $H' = 4.95 \pm .84$ for the tree buttress collection taken in Ecuador. Part of the within-habitat diversity differences are due to a difference in kind of organism: a large number of terrestrial frogs are present in Ecuador; there are no terrestrial frogs from the Thai site. Other differences are also operating, however. The number of lizard species are comparable in the two collections (8 from Thailand, 10 from Ecuador), yet the Brillouin diversity measure of the Ecuadorian lizard collection was $NH = 2.24$, higher than the diversity of the Thai lizard collection, $NH = 1.25$. The diversity differences are postulated to be due to different sizes of the total forest herpetofaunas of the two regions.

INTRODUCTION

The role of within- and/or among-habitat differences in explaining species diversity gradients has been a topic of much discussion based primarily on hypotheses with few direct data. The purpose of this paper is to examine the species diversities of amphibians and reptiles from apparently similar microhabitats from two different tropical forests. If the two forests are thought of as collections of microhabitats, certain of the microhabitats are equivalent, but the total number of microhabitats differ in each forest. If diversity differences are due solely to among-habitat differences, the diversities of similar microhabitats should be similar, but the total diversities would differ because of the unequal number of microhabitats in each forest. If within-habitat diversity differences are present in addition to among-habitat differences, the diversities of similar microhabitats will differ. Analysis should allow distinction of whether or not within-habitat differences are operating.

The tree buttress system was examined (Ecuador). The by either hoe (Thailand) or machete (Ecuador). The buttress system was examined to the point where the buttress originated from the trunk or to eye level in the few cases where the buttresses were taller than the collectors. When individual animals were captured, we recorded their exact location as well as the diameter of the tree at breast height. The sampling in Thailand was done in conjunction with random quadrat sampling, and covered the 10-month period of March–December 1969. The sampling in Ecuador was done specifically to determine the composition of the buttress herpetofauna and was done in June and July of 1971. In Ecuador, additional information on the extent of the buttresses and extent of litter lying between the buttresses was also recorded and an effort was made to sample the trees randomly. The data on tree diameters, buttress diameters, degree of leaf litter development, and occurrence of animals are available on request from the first author to anyone who wishes to utilize them. The Thai specimens are in the collections of the Field Museum of Natural History; the Ecuador specimens are in the collections of the Los Angeles County Museum of Natural History. Pielou's (1966) method and the Brillouin (1956) measure are used to determine the species diversities of the two collections. The Pielou method allows determination of a standard error and has already been used in analyzing quadrat data for amphibians and reptiles (Lloyd, Inger, and King 1968). In this case, each "quadrat" is a tree buttress that had amphibians or reptiles associated with it. Empty tree buttresses were not used.

METHODS AND MATERIALS

The tree buttress increases the interface of tree and ground. We examined this interface for reptiles and amphibians. The ground surface lying among buttress extensions was examined, the surface litter system for the tall, monolayered trees of tropical forests. The tree buttress system has evolved as a support system for the tall, monolayered trees of tropical forests. The buttress consists of flat, triangular plates

¹ Received June 16, 1972; accepted September 26, 1972.

which are confluent with the base of the tree trunk

and extend into the ground.

The buttresses are discrete units, easy to recognize

and sample. The similarity in appearance of the but-

tresses at the two sites prompted Heyer, who was

present at both sites, to initiate the comparative sam-

pling in Ecuador. As far as we know, there is no

information indicating whether or not amphibians

and reptiles utilize tree buttresses distinctively with

respect to either tree trunks or forest floor. However,

the tree buttress has distinctive features which would

be recognized by amphibians and reptiles: (1) crev-

ices often form where the buttress joins the trunk,

providing hiding places for lizards, (2) leaf litter is

deeper between the buttresses than on surrounding

ground, (3) caverns are often formed where the

buttress, ground, and trunk meet, providing hiding

places for surface animals.

In Thailand, the dry evergreen forest was sampled

over the entire year. An example illustrates that the

species are not restricted to tree buttresses, but are

commonly associated with them. Sixty-nine of 360

specimens (19%) of *Riopa bowringi*, the common-

est forest floor lizard, were found in tree buttresses.

SPECIES DIVERSITY RESULTS

The following 12 species of amphibians and rep-

tiles were collected from tree buttresses in Thailand.

Frogs
Family Microhylidae
Microhyla inornata - 1 individual

Lizards
Family Gekkonidae
Cyrtodactylus angulatus - 1

Peropus multatus - 2

Family Scincidae

Davewakium miritamae - 4

Leiopismsa reevesi - 8

Leiopismsa stamensis - 6

Mabuya multifasciata - 1

Riopa bowringi - 69

Sphenomorphus maculatus - 1

Snakes

Family Colubridae

Liopeltis scriptus - 2

Oligodon cyclurus - 1

Psammodrynastes pulverulentus - 1

Total Number of Individuals - 97

From tree buttresses in Ecuador we collected the following 33 species:

Caecilians

Family Caeciliidae

Caecilia sp. - 1 individual

Frogs

Family Bufonidae

Bufo typhonus - 9

Family Dendrobatiidae

Colostethus brunneus - 8

Phyllobates femoralis - 14

Phyllobates pictus - 1

Family Hyllidae

Hyla bokermanni - 2

Hyla funerea - 2

Hyla garbei - 1

Family Leptodactylidae

Eleutherodactylus acuminatus - 1

Eleutherodactylus brevicauda - 1

Eleutherodactylus conspicillatus - 7

Eleutherodactylus ockendeni - 1

Eleutherodactylus sulcatus - 2

Eleutherodactylus variabilis - 1

Eleutherodactylus sp. A - 11

Eleutherodactylus sp. B - 2

Ischnocnema guixensis - 1

Leptodactylus discodactylus - 1

Leptodactylus mystacens - 3

Leptodactylus wagneri - 5

Family Microhylidae

Chiasmocelis ventrimaculatus - 15

Hamptophryne boliviana - 10

Lizards

Family Gekkonidae

Gonatodes constrictus - 10

Pseudogonatodes sp. - 3

Family Iguanidae

Anolis sp. - 3

Erythrolates laticeps - 2

Family Teiidae

Iphisia elegans - 2

Leposoma parvifolium - 3

Ophiogommon abendrothi - 3

Ptychocheilus brevifrons - 2

Tupinambus nigropunctata - 1

Snakes

Family Colubridae

Rhadinaea brevirostris - 1

Total Number of Individuals - 143

sure is:

$$NH = \log_2 N! - \sum_{i=1}^s \log_2 n_i^i \quad (1)$$

where N is total individuals, n_i are the individuals in each species, and H is the mean diversity per individual in the collection itself. Once there is no change in diversity as sample size increases, the mean of successive values of h_k can be used as an estimator of H . Values for h_k are determined as

$$h_k = \frac{(N^k H^k - N^{k-1} H^{k-1})}{(N^k - N^{k-1})} \quad (2)$$

where the subscripts refer to the number of quadrats accumulated thus far in the process. For the Thai data, H' estimates were begun with the 50th buttress, for the Ecuadorian data, the 85th buttress (Fig. 1). We repeated the entire process 10 times, shuffling the data cards prior to each new estimation. The definitive estimates of H' and its standard error are the medians of the 10 runs. For the Thai data, $H' = 1.92 \pm .47$, for the Ecuador data, $H' = 4.95 \pm .84$.

For the Thai lizards, $NH = 1.25$, for the Ecuador lizards, $NH = 2.24$. The Ecuadorian lizard collection is clearly more diverse than the Thai lizard collection from tree buttresses.

DISCUSSION

The most obvious aspect of the difference in diversities between the Thai and Ecuadorian collections is the relative role that frogs play in each collection. In Thailand, only a single frog was taken from the buttress microhabitat. In Ecuador, 21 species of frogs were collected from the buttress microhabitat. This difference correlates with differences in anuran reproductive biology in the two regions. Two dominant frog groups of the buttress fauna from Ecuador have terrestrial breeding habits, or nearly so. Frogs of the genus *Eleutherodactylus* have direct development; eggs are laid in moist terrestrial situations, and small adults hatch from the eggs. Male frogs of the family Dendrobatiidae transport larvae from small puddle to small puddle in the forest; the frogs are terrestrial and do not require large rain pools or ponds for any portion of their life histories. Frogs with life cycles approaching direct development are exceptional in the anuran fauna of Southeast Asia. The herpetofaunas of Southeast Asia and South America have undergone essentially independent evolutionary histories and indicate that differences in species diversity may be due in part to phylogenetic differences when faunas are compared from differing biogeographic areas.

Even though the lizards of the two collections are distinct at the generic level, there are no obvious differences in biology such as is the case for the anurans. In fact, the gekkonids seemed to be ecological equivalents in both regions, inhabiting the buttress crevices on the trunks; and skinks, iguanids, and turtles were found in the buttress-litter-soil interface. A burrowing lizard was present in each area, *Davewakium mitrianae* in Thailand, *Ophiognomon abendrothii* in Ecuador. The differences in diversities of the two collections cannot be correlated with phylogenetic differences as such, but rather reflect differences due to the within-habitat component of species diversity. Karr (1971) has demonstrated that the among-habitat component of diversity does not account for the total diversity differences of temperate and tropical bird communities in the New World, but that within-habitat diversity differences are also operating. These conclusions corroborate those of Orans (1969) based on bird community comparisons from different vegetation formations in Central America. Both authors attribute the within-habitat avian diversity differences to the addition of new food categories in the more complex vegetation formations. Such an explanation is not likely for the within-habitat diversity differences noted here, especially

the Brillouin measures of diversity for the total collections are for the Thai data, $NH = 1.58$, for the Ecuador data, $NH = 3.92$. The diversities of the two collections are clearly different. One obvious difference between the collections is the importance of frogs in each. However, the number of lizard species is comparable in the two collections. The lizard subsets of the two collections were analyzed in the same manner as the total collections. Fig. 2 indicates the accumulative values of H . Estimates of H were begun with the 20th buttress for the Thai data (Fig. 2), at the point where H stabilizes. The average of 10 estimates of H and the associated standard error for the Thai lizard data is $H = 1.53 \pm .30$. Obviously the estimates of H have not leveled off for the Ecuadorian data (Fig. 2). The Brillouin measure provides a better measure in this case for the lizard components.

Fig. 1. Successive values of average diversity of amphibians and reptiles collected in tree buttresses from a moist tropical forest in Ecuador (curve A) and a dry evergreen forest in Thailand (curve B). The H was recalculated (Equation 1) following the addition of each buttress to the total collection, taking the buttresses in random order. Buttresses beyond the points indicated by the arrows were used to obtain h_2 of equation 2.





Fig. 2. Successive values of average diversity of lizards collected in tree buttresses from a moist tropical forest in Ecuador (Curve A) and a dry evergreen forest in Thailand (Curve B). See legend for Fig. 1 for further explanation.



gustics who allowed us to work at their base camp, Limoncocha, aided our work, especially the Don Johnsons and Bub Bormans.

Paul Liebelt, Pacific Lutheran University, programmed and ran the Pielou and Brillouin methods on the computer facilities provided by Pacific Lutheran University. William E. Duellman, University of Kansas, John D. Lynch, University of Nebraska, Jay M. Savage, University of Southern California, and Philip A. Silverstone and John W. Wright, both of the Los Angeles County Museum of Natural History, assisted in identifying some of the Ecuadorian material.

Robert F. Inger, Field Museum of Natural History, Dennis R. Paulson, University of Washington, and an anonymous reviewer read the manuscript and made helpful suggestions.

Fieldwork in Thailand was supported by National Science Foundation Grant GB7845X to Inger. Remaining support was from NSF Grant GB-27280 to Heyer.

LITERATURE CITED

- Brillouin, L. 1956. Science and information theory. Academic, New York.
- Holdridge, L. R. 1964. Life zone ecology. Tropical Science Center, San Jose, Costa Rica.
- Karr, J. R. 1971. Structure of avian communities in selected Panama and Illinois habitats. Ecol. Monogr. 41: 207-233.
- Lloyd, M., R. F. Inger, and F. W. King. 1968. On the diversity of reptile and amphibian species in a Bornean rain forest. Am. Nat. 102: 497-515.
- MacArthur, R. H., and E. O. Wilson. 1967. The theory of island biogeography. Princeton Univ. Press, Princeton.
- Orians, G. H. 1969. The number of bird species in some tropical forests. Ecology 50: 783-801.
- Pielou, E. C. 1966. The measurement of diversity in different types of biological collections. J. Theor. Biol. 13: 131-144.

ACKNOWLEDGMENTS

Many people aided our collecting efforts in Ecuador and Thailand. Especially helpful in Thailand were Prasert Lohavajjaya and Sukhum Pongsapattana of the Applied Scientific Research Corporation of Thailand. In Ecuador many people of the Summer Institute of Limoncocha, aided our work, especially the Don Johnsons and Bub Bormans.

Paul Liebelt, Pacific Lutheran University, programmed and ran the Pielou and Brillouin methods on the computer facilities provided by Pacific Lutheran University. William E. Duellman, University of Kansas, John D. Lynch, University of Nebraska, Jay M. Savage, University of Southern California, and Philip A. Silverstone and John W. Wright, both of the Los Angeles County Museum of Natural History, assisted in identifying some of the Ecuadorian material.

Robert F. Inger, Field Museum of Natural History, Dennis R. Paulson, University of Washington, and an anonymous reviewer read the manuscript and made helpful suggestions.

Fieldwork in Thailand was supported by National Science Foundation Grant GB7845X to Inger. Remaining support was from NSF Grant GB-27280 to Heyer.

Fieldwork in Thailand was supported by National Science Foundation Grant GB7845X to Inger. Remaining support was from NSF Grant GB-27280 to Heyer.